

Distribution of Phytochemicals and Bioactivity in Different Parts and Leaf Positions of *Stevia rebaudiana* (Bertoni) Bertoni

K.L.N.W. KUMARI¹, D.C. ABEYSINGHE¹ and R.M. DHARMADASA²

¹Department of Plantation Management, Faculty of Agriculture and Plantation Management, Wayamba University of Sri Lanka, Makandura, Gonawila (NWP), 60170, Sri Lanka

²Industrial Technology Institute, Bauddhaloka Mawatha, Colombo 07, 00700, Sri Lanka

ABSTRACT

Stevia rebaudiana (Bertoni) Bertoni (Asteraceae) is a small perennial herb which is widely grown for its sweet leaves and possesses 250-300 times the sweetness of sugar due to the presence of steviol glycosides (mainly stevioside and rebaudioside). Further, it is commonly known as candy leaf, sweet leaf and sugar leaf. However, studies on therapeutically important active components presence in stevia are scattered or lacking. Therefore, the present study was undertaken to determine the Total Phenolic Content (TPC), Total Flavonoid Content (TFC) and Total Antioxidant Capacity (TAC) of different parts of the plant and different leaf positions of *S. rebaudiana*. Leaf fresh weight, dry weight and leaf area at different leaf positions were recorded. The TAC, TPC and TFC were determined using Ferric Reducing Antioxidant Power Assay (FRAP), modified Folin-Ciocalteu colorimetric method and calorimetric method respectively. Leaf fresh weight and leaf dry weight were increased with the maturity. Further, all tested parts demonstrated the presence of TPC, TFC and TAC. The significantly higher TPC, TFC and TAC were reported in leaves than other parts of plant. The order of increase was leaf>flower>stem>branch>root. TPC, TFC and TAC of different leaf positions revealed that TAC, was decreased gradually from immature to mature leaf (1st leaf>2nd leaf>3rd leaf>4th leaf>5th leaf). Further, all tested phytochemicals (phenolics and flavonoids) and antioxidant capacity were significantly higher in extracts prepared from the first leaf. Strong significant correlations were observed between TAC and tested secondary metabolites (TPC, R²=0.85 and TFC, R²=0.85). The results of the present study are vital important in harvesting and quality control aspects of *S. rebaudiana*.

KEYWORDS: Antioxidant capacity, Flavonoids, Leaf position, Phenolics, Plant parts, *Stevia rebaudiana*

INTRODUCTION

Stevia rebaudiana (Bertoni) is a small perennial herb belonging to the family Asteraceae (Manoharan *et al.*, 2014). The plant is native to Paraguay and Brazil, and produces high potency low calorie sweeteners, stevioside and rebaudioside which are about 250-300 times sweeter than sucrose (Brandle *et al.*, 1998; Debnath, 2008; Katarzyna *et al.*, 2015; Rajesh *et al.*, 2010). These natural sweeteners are gaining very high popularity amongst all type of sweetener users as the most ideal substitute for sugar (Sheeja and Beena, 2015). Other than, its sweetening property stevia is also known for its therapeutic properties including antidiabetic (Mizutani and Tanaka, 2002), antimicrobial (Satishkumar *et al.*, 2008), antiviral (Kedik *et al.*, 2009), antifungal (Silva *et al.*, 2008), anti-hypertensive (Chan *et al.*, 1998; Lee *et al.*, 2001; Hsieh *et al.*, 2003), anti-hyperglycaemic (Jeppesen *et al.*, 2002; Benford *et al.*, 2006), anti-tumour (Satishkumar *et al.*, 2008), anti-human rota-virus activities (Das *et al.*, 1992; Takahashi *et al.*, 2001), anti-HIV (Takahashi *et al.*, 1998), hepatoprotective (Mohan and Robert., 2009) and immunomodulatory effects (Jaroslav *et al.*, 2006; Chatsudthipong and Muanprasat, 2009). Even though there are many information more

available on sweetening capacity, information on health benefits due to the presence of therapeutically active secondary metabolites are less available. Systematic studies on phytochemical distribution in different parts of the plant including leaves at different leaf positions are lacking. Therefore, present study was undertaken to determine the Total Antioxidant Capacity (TAC), Total Phenolic Content (TPC) and Total Flavonoid Content (TFC) of different parts (leaves, stems, roots, flowers and branches) of *S. rebaudiana* and different leaf positions of the plant (1st leaf, 2nd leaf, 3rd leaf, 4th leaf and 5th leaf).

MATERIALS AND METHODS

Location

The study was conducted in the laboratory of Physiology, Department of Plantation Management, Wayamba University of Sri Lanka, Makandura, Gonawila (NWP) from December 2015 to May 2016.

Chemicals and Regents

Folin-Ciocalteu reagent, Gallic acid, Rutin, 2, 4, 6-trypridyl-2-try-azine (TPTZ), 6-hydroxy-2, 5, 7, 8-tetramethy-chroman-2-carboxylic acid (Trolox) and ferric chloride (FeCl₃.6H₂O) were purchased from sigma

Aldrich Chemical Co. (St. Louis, Mo). All other chemicals used were of analytical grade.

Preparation of Samples

Samples of leaves, flowers, roots, stem parts and branches and also leaves from different leaf positions (1st, 2nd, 3rd, 4th and 5th leaf) of *Stevia rebaudiana* were collected from same aged, similarly maintained plants grown under same soil and climatic conditions (Low Country Intermediate Zone (IL_{1a}), at an elevation of 25 m above mean sea level with soil type of red yellow podosolic at experimental plots of Wayamba University of Sri Lanka, Makandura, Gonawila (NWP).

Extraction of Samples

Collected samples were cut into small pieces and air dried for three days at room temperature (28±2 °C). Then samples (Three samples from each category) were powdered using motor and pestle and sieved with 0.25 mm mesh. Powdered samples (0.1 g) were mixed with 5 mL of 80% methanol vortexed for 15 min. Then it was placed in a water bath at 60 °C for 40 min and vortex procedure was repeated at 10 min interval. After centrifugation at 4,000 rpm for 5 min, the supernatant was decanted into a 15 mL centrifuge tube and the remaining was re-extracted with 5 mL of 80% methanol. Supernatants were pooled and stored at -20 °C prior to analysis.

Total Phenolic Content (TPC)

Total phenolic content was quantified using a modified Folin-Ciocalteu method (Abeysinghe *et al.*, 2007). Briefly, 4 mL of distilled water and 0.5 mL of properly diluted leaf extract were mixed with 0.5 mL of 0.5 N Folin-Ciocalteu reagent (FCR) and allowed to react for 3 min. Then 1 mL saturated sodium carbonate solution was mixed and incubated in a water bath for 2 h at 30 °C. The absorbance was measured at 760 nm using UV visible spectrophotometer (Shimadzu, UV Mini 1240, Japan). Gallic acid was used as the standard and data were expressed as mg of Gallic Acid Equivalent (GAE) per gram of dry weights.

Quantification of Total Flavonoid (TFC)

Total flavonoid content (TFC) was determined by a colorimetric method, with slight modifications (Liu *et al.*, 2002). Briefly, 0.5 mL of the plant extract was diluted with 3.5 mL of distilled water. Then 0.3 mL of 5% NaNO₂ solution was added to the mixture. After 6 min, 0.3 mL of a 10% Al(NO₃)₃.6H₂O solution was added, and the mixture was allowed to stand for another 6 min. Then 2 mL of 2 M NaOH was added, and top up to 8 mL

with distilled water. After thoroughly mixing, the absorbance was measured at 510 nm using UV visible spectrophotometer (Shimadzu UV-160, Japan). Rutin was used as the standard and data were expressed as mg of Rutin Equivalent (RE) /g DW.

Determination of Total Antioxidant Capacity (TAC)

Total antioxidant capacity was determined using Ferric Reducing Antioxidant Power (FRAP) assay (Benzie and Strain, 1996) with slight modifications. Methanolic extract of sample (100 µL) was mixed with 900 µL of freshly prepared FRAP reagent of pH 3.6 containing 2.5 mL of 10 mol/L, 2,4,6-Tripyridyl-s-Triazine (TPTZ) solution in 40 mmol/L, HCl plus 2.5 mL of 20 mmol/L FeCl₃ and 25 mL of 300 mol/L acetate buffer. Absorbance was measured at 593 nm using the spectrophotometer (Shimadzu, UV Mini 1240, Japan) after incubating for 4 min. Trolox was used as the standard solution and TAC was expressed as mg Trolox Equivalents (TE) /g DW.

Determination of Leaf Area

Leaf area was measured using a leaf area meter (LI-3100C, USA). Samples were collected from 10 plants and ten leaves from each category (part from first to fifth leaves).

Determination of Fresh to Dry Weight Ratio

Samples were collected from 10 plants (part from first to fifth leaves) and fresh weight was measured then the samples were air dried. After three days dry weight of the samples were measured.

Statistical Analysis

To verify the statistical significance of all parameters the values of means and ±SD were calculated. Statistical comparison of mean values was performed by General Linear Model (GLM) of ANOVA followed by Turkey Multiple Range Test using SAS (SAS institute, 1999). The P values of less than 0.05 were adopted as statistically significant.

RESULTS AND DISCUSSION

In the present study attempts were made to determine economical yield, therapeutically important phytochemicals (phenolics and flavonoids) and Total Antioxidant Capacity (TAC) of different parts and different leaf positions of *S. rebaudiana*.

As demonstrated in Table 1, current study exhibited the presence of major phytochemical secondary metabolites (TPC and TFC) and TAC in all five main parts (leaves, stems

branches, flowers and roots) of *S. rebaudiana*. Significantly higher TAC, TPC and TFC were reported in leaf extracts. Order of increase was leaves>flowers>stem>branches>roots. The higher content of TAC in leaf and flower extract and TPC and TFC in leaf extract might be due to presence of high content of phenolics, flavonoids and variety of other pigments in the leaf. Systematic investigation of distribution of important phytochemicals is important for the quantitative estimation and locating of pharmacologically active chemical compounds (Sharanabasappa *et al.*, 2007). Presence of biologically active phytochemicals in all parts of the plant is important for the therapeutic activity of plant. Further, presence of higher content of investigated phytochemicals and antioxidant activity is in agreement with Abeysiri *et al.*, (2013), who investigated the phytochemical distribution of different parts of *Acmella oleraceae*. Moreover, the highest therapeutically active components presence in leaf and flower parts also similar with above investigation. Furthermore, higher content of TAC of leaf extracts are in agreement with Fernando, *et al.*, (2013) and Siriwardane *et al.*, (2013) who investigated the phytochemical distribution of *Withania somnifera*.

It is interesting to note that there were strong relationships between total antioxidant capacity and secondary metabolites; TPC ($R^2=0.85$) and TFC ($R^2=0.85$).

Economical yield of any plant depend on the physical and chemical yield of the plant. As shown in Figure 1 and Table 2, in the current study leaf fresh weight, leaf dry weight fresh to dry weight ratio, leaf area, TPC, TFC and TAC of different leaf positions were investigated. The results revealed that TAC was decreased

gradually from immature to mature leaf (1st leaf>2nd leaf>3rd leaf>4th leaf>5th leaf). Further, all tested phytochemicals (TPC and TFC) were significantly higher in extracts prepared from the first leaf (Bud).




Figure 1. Changes in area of leaves (A) and fresh weight, dry weight (B) at different leaf positions of *Stevia rebaudiana*

Table 1. Distribution of Total Antioxidant Capacity (TAC), Total Phenolic Content (TPC) and Total Flavonoid Content (TFC) in different parts of *Stevia rebaudiana*

| Plant part | TAC (mg TE / g DW) | TPC (mg GAE/ g DW) | TFC (mg RE / g DW) |
|------------|-------------------------|-------------------------|-------------------------|
| Leaves | 27.67±0.32 ^a | 12.44±0.02 ^a | 29.33±0.22 ^a |
| Flowers | 27.57±0.95 ^a | 11.99±0.03 ^b | 27.19±0.08 ^b |
| Roots | 03.45±0.30 ^d | 03.18±0.04 ^c | 01.95±0.30 ^c |
| Stems | 16.10±0.49 ^b | 08.49±0.01 ^c | 14.38±0.16 ^c |
| Branches | 10.76±0.33 ^c | 07.23±0.04 ^d | 13.76±0.08 ^d |

Means denoted by the same letters in a column represent non-significant differences ($p < 0.05$); TE-Trolox equivalent; GAE-Gallic acid equivalent; RE-Rutin equivalent; DW-Dry weight

Table 2. Total Antioxidant Capacity (TAC), Total Phenolic Content (TPC) and Total Flavonoid Content (TFC) and fresh to dry weight ratio of leaves at different positions of *Stevia rebaudiana*

| Leaf position | TAC (mg TE/g DW) | TFC (mg RE/g DW) | TPC (mg GAE/g DW) | Fresh to dry weight ratio |
|-----------------|-------------------------|-------------------------|-------------------------|---------------------------|
| 1 st | 34.24±0.78 ^a | 34.86±0.25 ^a | 13.64±0.04 ^a | 3.2:1 |
| 2 nd | 30.01±0.07 ^b | 23.29±0.14 ^d | 11.56±0.04 ^d | 2.9:1 |
| 3 rd | 28.10±0.68 ^c | 29.62±0.08 ^b | 12.67±0.02 ^b | 2.6:1 |
| 4 th | 24.80±0.44 ^d | 25.00±0.29 ^c | 11.23±0.06 ^c | 2.5:1 |
| 5 th | 21.15±0.49 ^c | 29.52±0.30 ^b | 11.77±0.05 ^c | 2.4:1 |

Means denoted by the same letters in a column represent non-significant differences ($p < 0.05$); TE-Trolox equivalent; GAE-Gallic acid equivalent; RE-Rutin equivalent; DW-Dry weight

CONCLUSIONS

Present study investigated Total Antioxidant Capacity (TAC), Total Phenolic Content (TPC) and Total Flavonoid Content (TFC) of different plant parts and different leaf positions of *Stevia rebaudiana*. Results indicated, all plant parts of *S. rebaudiana* possess TPC, TFC and TAC. The higher TPC, TFC and TAC were recorded leaf extracts than other parts of plant. Moreover, all tested secondary metabolite contents (TPC and TFC) and antioxidant capacity were reduced with the leaf maturity. Strong significant correlations were observed between TAC and tested secondary metabolites (TPC, $R^2=0.85$ and TFC, $R^2=0.85$).

ACKNOWLEDGEMENTS

Authors wish to express their gratitude to Mr. W.A.R. Wijesooriya, Technical Officer, Mr. H.M.A.S. Bandara and Mr. W.M.U.S. Bandara, Lab assistants and Mr. S.P.A. Karunaratne Field Supervisor, Department of Plantation Management, Wayamba university of Sri Lanka for their help during study period.

REFERENCES

- Abeyasinghe, D.C., Li, X., Sun, C., Zhang, W., Zhou, C. and Chen, K. (2007). Bioactive compounds and antioxidant capacities in different edible tissues of citrus fruit of four species. *Food Chemistry*, **104**, 1338–1344.
- Abeyisiri, G.R.P.I., Dharmadasa, R.M., Abeyasinghe, D.C. and Samarasinghe, K. (2013). Screening of phytochemical, physio-chemical and bioactivity of different parts of *Acmella oleraceae* Murr. (Asteraceae), a natural remedy for toothache. *Industrial crops and products*, **50**, 852-856.
- Benford, D.J., Dinovi, M. and Schlatter, J. (2006). Safety evaluation of certain food additives: Steviol Glycosides. WHO Food Additives Series (World Health Organization Joint FAO/WHO Expert Committee on Food Additives (JECFA), **54**, 140.
- Benzie, I.F. and Strain, J.J. (1996). The Ferric Reducing Ability of Plasma (FRAP) as a measure of "Antioxidant Power" the FRAP assay. *Analytical Biochemistry*, **239**, 70-76.
- Brandle, J. E., Starratt, A. N. and Gijzen, M. (1998). *Stevia rebaudiana*: Its agricultural, biological, and chemical properties. *Plant Science*, **78**, 527–536.
- Chan, P., Xu, D.Y., Liu, J.C., Chen, Y.J., Tomlinson, B., Huang, W.P. and Cheng, J.T. (1998). The effect of stevioside on blood pressure and plasma catecholamines in spontaneously hypertensive rats. *Life Science*, **63**, 1679-1684.
- Chatsudthipong, V. and Muanprasat, C. (2009). Stevioside and related compounds: Therapeutic benefits beyond sweetness. *Pharmacology Therapeutics*, **121**, 41-54.
- Das, S., Das, A.K., Murphy, R.A., Punwani, I.C., Nasution, M.P. and Kinghorn, A.D. (1992). Evaluation of the cariogenic potential of the intense natural sweeteners stevioside and rebaudioside A. *Caries Research*, **26**, 363–366.
- Debnath, M. (2008). Clonal propagation and antimicrobial activity of an endemic medicinal plant *Stevia rebaudiana*. *Journal of Medicinal Plants*, **2**, 45-51.
- Fernando, I.D.N.S., Abeyasinghe, D.C. and Dharmadasa, R.M. (2013). Determination of phenolic contents and antioxidant capacity of different parts of *Withania omnifera* (L) Dunal from three different growth stages. *Industrial crops and products*, **50**, 537-539.
- Hsieh, M.H., Chan, P., Sue, Y.M., Liu, J.C., Liang, T.H., Huang, T.Y., Tomlinson, B., Chow, M.S., Kao, P.F. and Chen, Y.J. (2003). Efficacy and tolerability of oral stevioside in patients with mild essential hypertension. *Clinical Therapeutics*, **25**, 2797-2808.
- Jaroslav, P., Barbora, H. and Tuulia, H. (2006). Characterization of *Stevia rebaudiana* by comprehensive two dimensional liquid chromatography time of flight mass spectrometry. *American Journal of Chromatography*, **1150**, 85-92.
- Jeppesen, P.B., Gregersen, S., Alstrup, K.K. and Hermansen, K. (2002). Stevio-side induces antihyperglycaemic, insulinotropic and glucagonostatic effects *in vivo*: studies in the diabetic Goto-Kakizaki (GK) rats. *Phytomedicine*, **9**, 9-14.
- Katarzyna, G.B., Tomasz, B., Zofia, N.L., Beata, A., Anna, J., Monika, K. and Kamila, R. (2015). Inhibitory effect of stevioside on calcium influx to produce anti-hypertension. *Planta Medicine*, **67**, 796-799.
- Kedik, S.A., Yarstsev, E.I. and Stanishevskaya, I.E. (2009). Antiviral activity of dried extracted of *Stevia*. *Journal of Pharmaceutical Chemistry*, **43**, 198-199.
- Lee, C.N., Wong, K., Liu, J., Chen, Y., Chen, J. and Chan, P. (2001). Inhibitory effect of stevioside on calcium influx to produce anti-hypertension. *Planta Medicine*, **67**, 796-799.
- Liu, M., Li, X.Q., Weber, C., Lee, C.Y., Brown, J. and Liu, R.H. (2002). Antioxidant and Anti-proliferative Activities of

- Raspberries. *Journal of Agriculture and Food Chemistry*, **50**, 2926-2930.
- Manoharan, R., Sivasangari, M., Kok, J., The, W.S. and Sarveisan, K. (2014). Studies on biochemical and medicinal properties of *stevia rebaudiana* grown *in vitro*. *International Journal of Ayurveda pharmaceutical*, **2** (5), 169-174.
- Mizutani, K. and Tanaka, O. (2002). Use of *Stevia rebaudiana* sweeteners in Japan. *Medicinal and Aromatic Plants*, **19**, 178-195.
- Mohan, K., Robert, J. (2009). Hepatoprotective effects of *Stevia rebaudiana* Bertoni leaf extract in CCl₄-induced liver injury in albino rats. *Plant Science*, **3**, 59-61.
- Rajesh, P., Rajesh, V.K. and Thambi, M.D. (2010). Effect of *Stevia rebaudiana* Bertoni ethanolic extract on anticancer activity of erlich's ascites carcinoma induced mice. *Journal of Current Biotica*, **3**, 549-554.
- Satishkumar, J., Sarvanan, M.M. and Seethalakshmi, I. (2008). *In-vitro* antimicrobial and antitumor activities of *Stevia rebaudiana* (Asteraceae) leaf extracts. *Journal of Pharmaceutical Research*, **7**, 1143-1149.
- Sharanabasappa, G.K., Mallikharjuna, P.B., Rajanna, L.N. and Seetharam, Y.N. (2007). Antibacterial and phytochemical evaluation of three medicinal plants. *Journal of natural products*. 327-334.
- Sheeja, R.R. and Beena, L. (2015). Phytochemical Screening of the Leaves of *Stevia rebaudiana*, Bertoni. *International Journal of Applied Sciences*, **4**, 344-347.
- Silva, P.A., Oliveira, D.F., Prado, N.R., Carvalho, D.A. and Carvalho, G.A. (2008). Evaluation of the antifungal activity by plant extracts against *Colletotrichum gloeosporioides* PENZ. *Ciencia Agrotecnologia*, **32**, 420-428.
- Siriwardane, A.S., Dharmadasa, R.M. and Kosala, S. (2013). Distribution of withaferin A, an anticancer potential agent, in different parts of two varieties *Withania somnifera* (L.). Dunal. Grown Sri Lanka. *Pakistan journal of Biological Sciences*, **16** (3), 141-144.
- Takahashi, K., Iwata, Y., Mori, S. and Shigeta, S. (1998). *In-vitro* anti- HIV activity of extract from *Stevia rebaudiana*. *Antiviral Research*. 37-59.
- Takahashi, K., Matsuda, M., Ohashi, K., Taniguchi, K., Nakagomi, O., Abe, Y., Mori, S., Sato, N., Okutani, K. and Shigeta, S. (2001). Analysis of anti-rotavirus activity of extract from *Stevia rebaudiana*. *Antiviral Research*, **49**, 15-24.