

Screening of Phytochemicals Distribution and Bioactivity of Different Parts and Leaf Positions of *Pimenta dioica* (L.) Merr (Myrtaceae) and Its Vegetative Propagation

P.B.C. PANAWALA¹, D.C. ABEYSINGHE¹ and R.M. DHARMADASA²

¹Department of Plantation Management, Faculty of Agriculture and Plantation Management, Wayamba University of Sri Lanka, Makandura, Gonawila (NWP), 60170, Sri Lanka

²Herbal Technology Section, Industrial Technology Institute, Baudhaloka Mawatha, Colombo 07, 00700, Sri Lanka

ABSTRACT

Pimenta dioica (L.) Merr. (Myrtaceae) is an evergreen aromatic spice widely used in perfumery, food and cosmetic industry in many parts of the world. Present study compared total antioxidant capacity (TAC), total phenolic content (TPC), total flavonoid content (TFC), leaf area (LA) and fresh to dry weight ratio of *Pimenta dioica* leaves at different maturity stages (1st, 2nd, 3rd, 4th and 5th leaf positions) and different plant parts (immature leaf, mature leaf and bark). Further an attempt has been made to investigate the optimum protocol for mass propagation of authentic, superior quality of *Pimenta dioica*. The TAC, TPC and TFC were determined using ferric reducing antioxidant power assay (FRAP), modified Folin-Ciocalteu method and calorimetric method respectively. Three types of cuttings (softwood, semi hardwood and hardwood) at two hormone level (with hormone and without hormone) were obtained from healthy mother plants and planted in poly bags filled with four different potting media (varying ratios of top soil: sand: compost (1:1:1, 1:1:2 and 1:2:1), pure sand only and sand+charcoal). Survival rate was recorded in monthly interval up to 3 months. Significantly higher TAC 562.38±9.42 (mg TE/g DW) and TPC 279.53±7.02 (mg GAE/g DW) were observed in leaf extract obtained from 1st leaf position. However the highest TFC 303.48±8.87 (mg RE/g DW) were observed in 5th leaf position. According to phytochemical distribution pattern, significantly higher TAC (619.84±11.98 mg TE/g DW), TPC (267.53±5.03 mg GAE/g DW) and TFC (305.48±8.87 mg RE/g DW) were observed in extracts obtained from bud region. Higher survival rate was observed in sand+charcoal potting media in soft wood cuttings. It is thus suggested to use softwood as planting material and sand+charcoal as potting medium for propagation of *Pimenta dioica* in commercial scale cultivation.

KEYWORDS: Antioxidant capacity, Flavonoids, Phenolics, *Pimenta dioica*, Vegetative propagation

INTRODUCTION

Pimenta dioica (L.) Merr. (Myrtaceae) is well known for its characteristic aroma, therapeutic and culinary qualities which resembles to aroma and flavor of clove, nutmeg cinnamon and hence it is called allspice (Weiss, 2002). Plant is known as, Jamaica pepper, Myrtle pepper and new spice in English (Jirovetz *et al.*, 2007). Further, *P. dioica* is small (7-10 m) evergreen tree which is cultivated in Jamaica, Cuba, Haiti, Brazil, Central America, West Indies, Venezuela, Mexico, Honduras, Guatemala and Grenada (Dharmadasa *et al.*, 2015). Leaves and seeds are widely used as commercial items in traditional medicine for the treatment of diverse range of ailments including flatulence, diarrhea, neuralgia, rheumatism and digestive problems (Anon, 1969). Moreover, it is valued for its secondary metabolites like essential oil with compositions, oleoresin which is highly utilized in food industry, perfumery and cosmetic industry. These diverse ranges of utilities may be due to presence of therapeutically active secondary metabolites in both seeds and leaves of *P. dioica*. On the other hand, once important

plant source found, its commercial planting material production is vital important. Allspice is conventionally propagated through seeds, but the seeds lost their viability soon after harvest (Rema *et al.*, 2008).

Allspice being a polygamodioecous tree, identification of the functional male and female trees till they flower is difficult and hence, clonal propagation is essential to obtain uniformly high yielding trees (Rema *et al.*, 1997). The substances, traditionally referred to as secondary metabolites or phytochemicals, often are differentially distributed among limited taxonomic groups within the plant kingdom (Croteau *et al.*, 2000).

Phytochemicals are chemical compounds that occur naturally in plants. Some are responsible for colour and other organoleptic properties. The most important of these bioactive constituents of plants are alkaloids, tannins, flavonoids and phenolic compounds (Hill, 1952). Secondary metabolites are those metabolites which are not functioning in growth (although they may have survival function). Many of these compounds now have been shown to have important adaptive significance

in protection against herbivory and microbial infection, as attractants for pollinators and seed dispersing animals and as allelopathic agents (allele chemicals that influence competition among plant species). Total phenolic content (TPC), total flavonoid content (TFC) and total antioxidant capacity (TAC) have been renowned as therapeutically active substances which regulate important physiological actions on the human body (Chandur *et al.*, 2011). Since antioxidant function as free radical scavengers and quenchers of singlet oxygen formation, investigation of natural antioxidants has greatly increased in recent years (Kaufman *et al.*, 1999). Out of the commonly used spice crops, allspice is distinguished due to its multi aroma and flavor characteristics (Jirovetz *et al.*, 2007).

Although there are scattered information available on different phytochemicals present in this plant, comparative study on the most important maturity stages are yet to be studied. Therefore, present study mainly having understood this enormous necessity, present study was undertaken to develop a simple, cost effective vegetative propagation protocol for mass propagation of *Pimenta dioica*.

MATERIALS AND METHODS

Location

Experiment was carried out in the experimental plots and the laboratory of the Department of Plantation Management, Wayamba University of Sri Lanka, Makandura, Gonawila (NWP) from December 2015 to June 2016. The experimental site was situated in the Low Country Intermediate-Zone (IL_{1a}), at an elevation of 25 m above mean sea level (Panabokke, 1996). The average annual temperature, relative humidity and solar intensity of the experimental area are 31°C, 78% and 85 kilo lux respectively.

Experimental Design and Treatments

Experimental design was a three factor factorial in randomized complete block design (RCBD). Three factors were potting medium (varying ratios soil: sand: compost 1:1:1, 1:1:2 and 1:2:1, pure sand only, and sand+charcoal), hormonal effect (with and without hormone) and type of cuttings (softwood, semi hardwood and hardwood). There were thirty (30) treatment combinations with ten (10) replicates. Plants (300) were treated with each treatment combinations. Each single pot was contained a cutting.

Planting Materials

Softwood, semi-hardwood and hardwood cuttings were obtained from 40 years old, well

maintained, healthy *Pimenta dioica* mother plants cultivated at Kundasale (GPS latitude 7.28291, GPS Longitude 80.6726), mid country intermediate zone (IM_{2b}), Kandy, Sri Lanka.

Preparation of Stem Cuttings

Vigorously grown healthy shoots were selected from mother plants to obtain cuttings. Three different maturity stages (softwood, semi-hardwood and hardwood) were obtained from selected shoots and cuttings were kept in a bucket of water until planting.

Preparation of Bags with Potting Medium

Gauge 150, transparent polythene bags (10 × 15 cm) were filled with three different potting media as follows. Medium 1- topsoil: sand:compost (1:1:1), medium 2- topsoil:sand:compost (1:1:2), medium 3- topsoil:sand:compost (1:2:1), medium 4- pure sand and medium 5- sand+charcoal. The top soil was taken from the university field where the land suitability class was low country intermediate-zone (IL_{1a}) in Sri Lanka. The compost fertilizer was provided from the research station near the university premises which consisted of poultry manure, cow dung, plant residuals and house litter in equal ratios.

Planting and Aftercare Operations

Three different maturity stages with three different hormone treatments (Hormone-Rapid root, Indole-3-butric acid and without hormone) were planted separately in the poly bags having five different potting media arranged in 30 treatment combinations. Immediately after planting polythene propagator (gauge 250, transparent) was established and shade was provided by coconut (*Cocos nucifera*) leaves. Propagator was maintained with similar conditions to minimize environmental variation. Pressure of hydrogen (pH) was maintained at seven for better nutrient absorption.

Measurement of Leaf Area

Leaf area was measured by a leaf area meter (LI-3100C, USA).

Extraction of Phytochemicals

Leaves were cut into pieces and air dried for three days at room temperature (28±2 °C). Then samples (three samples from each category) were powdered using motor and pestle and sieved with 0.25 mm mesh.

Powdered samples (0.1 g) were mixed with 5 mL of 80% methanol vortexed for 15 min. Then it was placed in a water bath at 60 °C for 40 min and vortex procedure was repeated in 10 min interval. After centrifugation at 4,000

rpm for 5 min, the supernatant was decanted into a 15 mL centrifuge tube and the remaining was re-extracted with 5 mL of 80% methanol.

Chemicals and Regents

6-hydroxy-2,5,7,8-tetramethy-chroman-2-carboxylic acid (Trolox), Gallic acid, Rutin, 2,4,6-tripyridyl-2-try-azine (TPTZ), Folin-Ciocalteu reagent and ferric chloride ($\text{FeCl}_3 \cdot 6\text{H}_2\text{O}$) were purchased from Sigma Aldrich Chemical Co. (St. Louis, Mo) and all other chemicals used were analytical grade.

The Total Phenolic Content (TPC)

The total phenolic content was quantified using modified Folin-Ciocalteu method (Abeysinghe *et al.*, 2007). Briefly, 4 mL of distilled water and 0.5 mL of properly diluted leaf extract were mixed with 0.5 mL of 0.5 N Folin-Ciocalteu reagent (FCR) and allowed to react for 3 min. Then 1 mL saturated sodium carbonate solution was mixed and incubated in a water bath for 2 h at 30 °C. The absorbance was measured at 760 nm using UV visible spectrophotometer (Shimadzu UV Mini 1240 Japan). Gallic acid was used as the standard and data were expressed as mg of Gallic Acid Equivalent (GAE) /g DW.

Quantification of Total Flavonoid (TFC)

Total flavonoid content (TFC) was determined by a colourimetric method, with slight modifications (Liu *et al.*, 2002). Briefly, 0.5 mL of the plant extract was diluted with 3.5 mL of distilled water. Then 0.3 mL of a 5% NaNO_2 solution was added to the mixture. After 6 min, 0.3 mL of a 10% $\text{Al}(\text{NO}_3)_3 \cdot 6\text{H}_2\text{O}$ solution was added, and the mixture was allowed to stand for another 6 min. Then 2 mL of 2 M NaOH was added, and top up to 8 mL with distilled water. After thoroughly mixing, the absorbance was measured at 510 nm using UV-visible spectrophotometer (Shimadzu UV Mini 1240 Japan). Rutin was used as the standard and data were expressed as mg of rutin equivalent (RE) /g DW.

Determination of Total Antioxidant Capacity (TAC)

Total antioxidant capacity was determined using Ferric Reducing Antioxidant Power (FRAP) assay (Benzie and Strain, 1996). Methanolic leaf extract (100 μL) was mixed with 900 μL of freshly prepared FRAP reagent of pH 3.6 containing 2.5 mL of 10 mol/L TPTZ solution in 40 mmol/L HCl plus 2.5 mL of 20 mmol/L FeCl_3 and 25 mL of 300 mol/L acetate buffer. Absorbance was measured at 593 nm using the spectrophotometer (Shimadzu, UV Mini 1240, Japan) after incubating for 4 min.

The trolox was used as the standard solution and TAC was expressed as mg Trolox Equivalents (TE) /g DW.

Statistical Analysis

Statistical comparison of mean values was performed by general linear model (GLM) of ANOVA followed by Turkey multiple range test of the SAS software package (SAS Institute, 1999).

RESULTS AND DISCUSSION

Spices are products derived from different parts of the aromatic plants *i.e.* leaves, roots, stem, branches, flowers, seeds, stigmas and styles or the entire plant tops which are heavily used by ethnic groups for their flavor, colour, aroma and preservation for foods or beverage worldwide (Anon, 2005).

The secondary metabolites (phenolics and flavonoids) and bioactivity play an important role in plant therapeutic value of medicinal plants. These secondary metabolites support mainly for the natural defense system, and mainly depend on the development stage of plants (Menikovic *et al.*, 2000; Fernando *et al.*, 2013). In the present study attempts were made to compare the variation of total antioxidant capacity, total phenolic content and total flavonoid content of leaves of *Pimenta dioica* at different maturity stages (Table 1). Total antioxidant capacity, was decreased with the maturity of leaves and TAC of 1st, 2nd, 3rd leaf positions were significantly different from 4th and 5th leaf positions.

As demonstrated in Table 1, Total phenolic content and total flavonoid content were decreased up to 3rd leaf position. However, they were significantly increased at 5th leaf position. Meanwhile leaf area was increased up to 4th leaf position and decreased at 5th leaf position (Table 1). Presence of higher content of TPC, TFC and TAC in immature leaves and bud, might be due to the less cellulose and pectin in immature leaves and the active components present in tender parts are more soluble than those present in mature parts. Observed high TAC, TPC and TFC in immature leaf and bud, extracts are in agreement with previous studies (Dharmadasa *et al.*, 2015).

Who investigated the higher TAC, TPC and TFC in immature and mature *Pimenta dioica* leaves and who investigated immature and mature tea (*Camellia sinensis*) leaves (Ming Yuan *et al.*, 2014).

Table 2 demonstrates the distribution of total antioxidant capacity, total phenolic content, total flavonoid content in different parts (leaf bud, immature leaf, mature leaf and bark) of *P. dioica*.

Table 1. Total antioxidant capacity (TAC), total phenolic content (TPC), total flavonoid content (TFC), leaf area and dry weight:Fresh weight ratio of leaves of *Pimenta dioica* at different leaf positions

| Leaf positions | TAC (mg TE/g DW) | TPC (mg GAE/g DW) | TFC (mg RE/g DW) | Leaf area (cm ²) | Dry weight:Fresh weight ratio |
|-----------------|---------------------------|--------------------------|--------------------------|------------------------------|-------------------------------|
| 1 st | 562.38±9.42 ^a | 279.53±7.02 ^a | 296.43±1.89 ^a | 0.6685±0.0002 | 37.80:100 |
| 2 nd | 552.36±1.63 ^a | 206.87±2.31 ^b | 291.90±1.49 ^a | 13.8515±0.0090 | 41.91:100 |
| 3 rd | 546.12±8.69 ^a | 154.20±8.72 ^c | 259.52±6.44 ^b | 39.209±0.113 | 49.51:100 |
| 4 th | 522.82±2.86 ^b | 154.20±5.29 ^c | 269.53±1.80 ^b | 50.4705±0.00305 | 52.12:100 |
| 5 th | 501.41±11.04 ^c | 152.87±1.15 ^c | 303.48±8.87 ^a | 42.082±0.01033 | 53.59:100 |

Numbers within a column followed by different letter are significantly different ($P < 0.05$), GAE- Gallic acid equivalent, TE- Trolox equivalent, RE- Rutin equivalent

Table 2, Total antioxidant capacity (TAC), total phenolic content (TPC), total flavonoid content (TFC) of *Pimenta dioica* at different plant parts

| Parts | TAC (mg TE/g DW) | TPC (mg GAE/g DW) | TFC (mg RE/g DW) |
|---------------|---------------------------|--------------------------|--------------------------|
| Bud | 619.84±11.98 ^a | 267.53±5.03 ^a | 305.48±8.87 ^a |
| Immature leaf | 395.99±11.10 ^b | 192.20±8.00 ^b | 225.48±5.95 ^c |
| Mature leaf | 325.80±10.29 ^c | 131.53±3.06 ^c | 221.90±3.30 ^c |
| Bark | 317.67±13.14 ^c | 143.53±8.33 ^c | 245.71±1.43 ^b |

Numbers within a column followed by different letter are significantly different ($P < 0.05$); GAE- Gallic acid equivalent, TE- Trolox equivalent, RE- Rutin Equivalent

The results clearly demonstrated that the highest TAC, TPC and TFC were contained in leaf bud. Order of decrease of total antioxidant capacity was leaf bud>immature leaf>mature leaf>bark.

The highest TAC was observed in bud (619.84±11.98 mg TE/g DW) and the lowest TAC was observed in bark (317.67±13.14 mg TE/g DW). As shown in Table 2, TFC, bud portion significantly different from bark portion, and bark showed higher significant difference than immature and mature leaf.

The highest TPC and TFC were observed in bud (267.53±5.03 mg GAE/g DW and 305.48±8.87 mg RE/g DW respectively). The lowest TPC and TFC were observed in mature leaf (131.53±3.06 mg GAE/g DW and 221.90±3.30 mg RE/g DW respectively). The correlation value between TAC and TPC was $R^2=0.9462$.

Effect of different potting media, cutting types and hormonal levels were evaluated and found that the higher survival rate was observed in sand+charcoal potting medium in softwood cuttings. It is thus suggested to use softwood as planting material and sand+charcoal as potting medium for propagation of *Pimenta dioica* in commercial scale cultivation.

CONCLUSIONS

Present study investigated phytochemicals distribution and bioactivity in different parts and different leaf positions and vegetative propagation protocol for mass propagation of *Pimenta dioica*. All tested bioactive compounds were distributed in all tested parts and the highest bioactive compounds were reported

from leaf or leaf bud positions. Further, present study clearly indicated that leaf can be used as raw materials for industrial purposes instead of fruits. Moreover, sand+charcoal potting medium in softwood cuttings could be used for propagation of *Pimenta dioica* in commercial scale.

ACKNOWLEDGEMENTS

Authors wish to express their gratitude to Mr. K.H.M.I. Karunarathna, instructor Information and communication Technology (ICT) Center, for great support with the statistical analysis. All non-academic members in Department of Plantation Management, Wayamba University of Sri Lanka and Mr. Nawarathna, home garden owner, Kundasale, Kandy, Sri Lanka, for their valuable assistance given to conduct this research study successfully.

REFERENCES

- Abeyasinghe, D.C., Li, X., Sun, C., Zhang, W., Zhou, C. and Chen, K. (2007). Bioactive compounds and antioxidant capacities in different edible tissues of citrus fruit of four species. *Food Chemistry*, **104**, 1338–1344.
- Anon. (1969). Wealth of India (Raw Materials). Publications Information Division, CSIR Publications. New Delhi, **8**, 58-59.
- Anon. (2005). Herbs, spices and essential oil post-harvest operations in developing countries. UNIDO Vienna, Austria.
- Benzie, I.F.F. and Strain, J.J. (1996). Ferric reducing ability of plasma (FRAP) as a measure of antioxidant power. The FRAP

- assay. *Analytical Biochemistry*, **239**, 70-76.
- Chandur, U., Shashidhar, S., Chandrasekar, S. B., Bhanumathy, M. and Midhun, T. (2011). Phytochemical evaluation and anti-arthritic activity of root of *Saussurea lappa*. *Pharmacologia*, **2**, 265-267.
- Croteau, R., Kutchan, I.M. and Lewis, N.G. (2000). Natural Products (Secondary Metabolites). *American Society of Plant Physiologist*, **24**.
- Dharmadasa, R.M., Abeysinghe, D.C., Dissanayake, D.M.N., Abeywardhana, K.W. and Fernando, N.S. (2015). Leaf essential oil composition, antioxidant activity, Total phenolic content, Total flavonoid content of *Pimenta dioica*. *Universal Journal of Agriculture Research*, **3** (2), 49-52.
- Fernando, D.N.S., Abeysinghe, D.C. and Dharmadasa, R.M. (2013). Phenolic content and antioxidant capacity of different parts of *Withania somnifera* (L) Dunal from different growth stages. In: proceedings of 12th Agricultural Research Symposium, 30-31 May, 2013. Wayamba university of Sri Lanka. 160-164.
- Hill, A.F. (1952). *Useful plants and plant products*. 2nd ed. Mc Garw-Hill book company Inc, New York.
- Jirovetz, L., Buchbauer, G., Stoilova, I., Krastanov, A., Stoyanova, A. and Schmidt, E. (2007). Spice plants, chemical composition and antioxidant properties of *Pimenta dioica* (L.) Merr. Leaf and oil from Jamaica. *Ernährung Nutrition*, **31**, 55-63.
- Kaufman, P.B., Cseke, L.J., Warber, S., Duke, J.A. and Briemann, H.L. (1999). *Natural Products from Plants*, CRC Press, Boca Raton, USA.
- Liu, M., Li, X.Q., Weber, C., Lee, C.Y., Brown, J. and Liu, R.H. (2002). Antioxidant and anti-proliferative activities of raspberries. *Journal of Agriculture and Food Chemistry*, **50** (10), 2926-2930.
- Menikovic, N., Savikin-Fodulovic, K. and Savin, K. (2000). Chemical compositions and seasonal variation in the amount of secondary compounds in *Gentiana lutea* leaves and flower. *Planta Medica*, **66**, 178-180.
- Ming, Y., Xuejing, J., Chunbang, D., Shu, Y., Zhongwei, Z. and Yanger, C. (2014). Comparative Studies on Bioactive Constituents in Hawk Tea Infusions with Different Maturity Degree and Their Antioxidant Activities. *The Scientific World Journal*, 1-7.
- Panabokke, C.R. (1996). Soils and Agro-ecological environments of Sri Lanka. Natural Resources Series No. 2. *Natural Resource, Energy and Science Authority of Sri Lanka, Colombo*, 219.
- Rema, J., Krishnamoorthy, B., Mathew, P.A. (1997). Vegetative propagation of major tree spices. *Journal of Spices and Aromatic Crops*, **6** (2), 57-405.
- Rema, J., Mathew, P.A. and Krishnamoorthy, B. (2008). Vegetative propagation of allspice. *Indian Journal of Horticulture*. 65-68.
- Weiss, E.A. (2002). *Spice Crops* CAB Publishing USA. 104-120.